


Alternative approach to fit irregular corneas

Julián Espinosa Tomás



Universitat d'Alacant
Universidad de Alicante



Universitat d'Alacant
Universidad de Alicante

Alternative approach to fit irregular corneas

Collaborators

Jorge Pérez


David Mas

Carmen Vázquez

Carlos Illueca

Funding


This work has been partially financed by the project
GV/2009/002 of the Conselleria d'Educació of the
Generalitat Valenciana





Background

Scheimpflug cameras and topographers based in Placido rings permit accurate estimation of corneal surfaces





Background

Corneal height data → Model → Wavefront analysis

☐ Zonal approach.

B-Splines

Well suited for fitting complex-shaped surfaces


Not related with aberrations

☐ Modal approach.

Zernike polynomials

Direct relation with Seidel aberrations

Not precise when describing highly irregular corneas





Objective


COMBINATION

ZONAL + MODAL approaches

Zonal Zernike fitting of corneal height data

Zernike coefficients computed in overlapping local areas

- ❑ Diminishing the influence of smooth areas over irregular zones and vice versa.
- ❑ Limiting the influence of the peripheral irregularities over the central corneal area, thus giving accurate reconstruction of the central optical zone.

























Modal approach. Zernike polynomials

$$W(x_u, y_v) \approx \sum_{j=0}^{p-1} c_j Z_j(x_u, y_v)$$

Least-squares method

$$C = (Z^T Z)^{-1} Z^T W$$





Modal approach. Zernike polynomials

$$W(x_u, y_v) \approx \sum_{j=0}^{p-1} c_j Z_j(x_u, y_v)$$

Least-squares method

$$C = (Z^T Z)^{-1} Z^T W$$

$$C = \begin{bmatrix} c_0 \\ c_1 \\ \dots \\ c_{p-2} \\ c_{p-1} \end{bmatrix} \quad Z = \begin{bmatrix} \sum_{j=0}^{p-1} Z_j(x_1, y_1) & \dots & \sum_{j=0}^{p-1} Z_j(x_n, y_1) \\ \dots & \dots & \dots \\ \sum_{j=0}^{p-1} Z_j(x_1, y_n) & \dots & \sum_{j=0}^{p-1} Z_j(x_n, y_n) \end{bmatrix}$$


$$W = \begin{bmatrix} W(x_1, y_1) & W(x_2, y_1) & \dots & W(x_{n-1}, y_1) & W(x_n, y_1) \\ W(x_1, y_2) & W(x_2, y_2) & \dots & W(x_{n-1}, y_2) & W(x_n, y_2) \\ \dots & \dots & \dots & \dots & \dots \\ W(x_1, y_{n-1}) & W(x_2, y_{n-1}) & \dots & W(x_{n-1}, y_{n-1}) & W(x_n, y_{n-1}) \\ W(x_1, y_n) & W(x_2, y_n) & \dots & W(x_{n-1}, y_n) & W(x_n, y_n) \end{bmatrix}$$



Zonal Zernike fitting

Auxiliary $a \times b$ matrices of size $N \times N$, unmasking $M \times M$ zone


$$W_{a,b}(u,v) \simeq \begin{cases} \sum_{j=0}^{p-1} c_j^{(a,b)} Z_j(\rho_{(u,v)}, \theta_{(u,v)}); \\ 0; \end{cases}$$






Zonal Zernike fitting

Modal



(N×N)

Modal+Zonal



(M×M)

One point of the surface may belong to different local regions




Zonal Zernike fitting

Reconstructed surface:
Mean at each point


$$L(u, v) = \frac{\sum_{a=1}^{N-(M-1)} \sum_{b=1}^{N-(M-1)} W_{a,b}(u, v)}{\sum_{a=1}^{N-(M-1)} \sum_{b=1}^{N-(M-1)} O_{a,b}(u, v)}$$

$$\sum_{a=1}^{N-(M-1)} \sum_{b=1}^{N-(M-1)} O_{a,b}(u, v)$$



Points in the central
zone are evaluated
 M^2 times

$M=21$ px \rightarrow 441 times






Zonal Zernike fitting

Number of elements in the central zone

$$f_N(\mathbf{M}) = \mathbf{M}^2 [\mathbf{N} - 2(\mathbf{M} - 1)]^2$$



From the derivative:

$$\mathbf{M}_{\text{opt}} = \text{round}\left(\frac{\mathbf{N} + 2}{4}\right)$$



Results

□ Height data analysis of surfaces.

□ Irregular surface

□ Keratoconus from Pentacam


Root mean square deviation (RMSD)

w_i = original surface data

g_i = reconstructed surface data

$$RMSD = \sqrt{\frac{\sum_{i=1}^{N'} (w_i - g_i)^2}{N'}}$$


□ Validate M_{opt} . Two masks of different sizes





Irregular surface: Sphere+Franke's function

$$fr(x, y) = \frac{3}{4} \exp \left[\frac{-((9x-2)^2 + (9y-2)^2)}{4} \right] + \frac{3}{4} \exp \left[- \left(\frac{(9x+1)^2}{49} + \frac{(9y+1)^2}{10} \right) \right] \\ + \frac{1}{2} \exp \left[- \left((9x-7)^2 + \frac{(9y-3)^2}{4} \right) \right] + \frac{1}{5} \exp \left[-((9x-4)^2 + (9y-7)^2) \right]$$






RMSD Sphere+Franke's function

Pupil diameter = 4 mm


$N=41 \rightarrow M_{\text{opt}}=11$






Differences Sphere+Franke's function


Modal: difference with 120 Zernike polynomials



Zonal mask 21 px: difference with 96 Zernike polynomials




Zonal mask 11 px: difference with 43 Zernike polynomials





Real irregular surface: Keratoconus

Keratoconus height data - sphere






RMSD keratoconus

Pupil diameter = 4 mm


$N=41 \rightarrow M_{\text{opt}}=11$






Differences keratoconus


Modal: difference with 178 Zernike polynomials



Zonal mask 21 px: difference with 196 Zernike polynomials



Zonal mask 11 px: difference with 217 Zernike polynomials






Conclusions

- ❑ Better results than modal fit for low order Zernike polynomials.
- ❑ The central surface part is better evaluated than the outer parts, since calculation is more intensive in this zone and not affected by peripheral irregularities
- ❑ Diminishes the influence of smooth areas over irregular zones and vice versa



Universitat d'Alacant
Universidad de Alicante



OCIVIS

<http://web.ua.es/es/ocivis/>